

## LIST OF FIGURES

<b><u>Figure</u></b>	<b><u>Description</u></b>	<b><u>Page</u></b>
2.1	Tandem solid phase/liquid trap	9
3.1	Chemical Structures for Lovastatin (a) and 17 $\alpha$ -Hydroxyprogesterone (b)	23
3.2	Effect of Methanol-Modifier Concentration on Lovastatin Recoveries (n=3) From In-House Prepared Tablet Powder Mixture	28
3.3	Effect of Additive Type on Lovastatin Recoveries (n=3) From In-House Prepared Tablet Powder Mixture	30
3.4	Effect of Methanol-Modifier Concentration on Lovastatin Recoveries (n=3) From In-House Prepared Tablet Powder Mixture	33
3.5	Subcritical Fluid Extraction (n=1) of Lovastatin From MEVACOR® Tablets at Various Additive/Modifier Concentrations	35
3.6	Subcritical Fluid Extraction (n=1) of Lovastatin From MEVACOR® Tablets at Various Modifier Concentrations With and Without IPA	36
3.7	Subcritical Fluid Extraction (n=1) of Lovastatin From MEVACOR® Tablets at Various Modifier Concentrations With and Without IPA	37
4.1	Chemical Structure of Triphenylphosphinetrisulfonate, Sodium Salt	53
4.2	Average Percent Recovery (n=3) TPPTS vs Methods Investigated	55
4.3	Significant Main Effects on Recovery	59

<b><u>Figure</u></b>	<b><u>Description</u></b>	<b><u>Page</u></b>
5.1	Chemical Structure of Pseudoephedrine Hydrochloride	64
5.2	Average Percent Recoveries (n=3) of Pseudoephedrine and Pseudoephedrine with 1-Heptanesulfonic Acid, Sodium Salt from a Non-Dried Drug Spiked-Sand Surface	86
5.3	Average Percent Recoveries (n=3) of Pseudoephedrine and Pseudoephedrine with 1-Heptanesulfonic Acid, Sodium Salt from a Dried Drug Spiked-Sand Surface	88
5.4	Average Percent Recoveries (n=3) of Pseudoephedrine with and without Various Ion-Pairing Additives at Various Concentrations from a Drug Spiked-Sand Surface	91
5.5	Average Percent Recoveries (n=3) of Pseudoephedrine with Various Modifiers and Ion-Pairing Additives from a Drug Spiked-Sand Surface	95
5.6	Average Percent Recoveries (n=3) of Pseudoephedrine from a Drug Spiked-Sand Surface. Effect of Addition of Acid and Base.	97
5.7	Average Percent Recovery (n=3) of Pseudoephedrine from Suphedrine Tablets as a Function of In-Cell Modifier Composition with Pure CO <sub>2</sub>	101
5.8	Average Percent Recovery (n=3) of Pseudoephedrine from Suphedrine Tablets as a Function of In-Cell Modifier Composition with 10% Methanol-Modified CO <sub>2</sub>	103
5.9	Average Percent Recovery (n=3) of Pseudoephedrine from Suphedrine Tablets as a Function of In-Cell Modifier Composition with 10% (1% H <sub>2</sub> O) Methanol-Modified CO <sub>2</sub>	105
5.10	Infrared Spectrum of Pseudoephedrine Hydrochloride Standard	109
5.11	Infrared Spectrum of a Suphedrine Tablet Extract With 10% Methanol-Modified CO <sub>2</sub>	110

<b><u>Figure</u></b>	<b><u>Description</u></b>	<b><u>Page</u></b>
5.12	Infrared Spectrum of a Suphedrine Tablet Extract With 10% (1% H <sub>2</sub> O) Methanol-Modified CO <sub>2</sub>	111
6.1	Chemical Structures of Phospholipids Investigated	119
6.2	Chromatogram A of 5 phospholipids	123
6.3	Chromatogram B of 3 phospholipids	125
6.4	Chromatogram C of 3 phospholipids	126
6.5	Chromatogram D of 5 phospholipids	128
6.6	Chromatogram E of 5 phospholipids	130
6.7	Chromatogram F of 5 phospholipids	131
6.8	Reproducibility comparison of 2 detectors	136

## LIST OF SCHEMES

<b><u>Scheme</u></b>	<b><u>Description</u></b>	<b><u>Page</u></b>
5.1	Phase I General Experimental Scheme	69
5.2	Phase II General Experimental Scheme	71
5.3	Phase III General Experimental Scheme	72
5.4	Phase IV General Experimental Scheme	73

## LIST OF TABLES

<b><u>Table</u></b>	<b><u>Description</u></b>	<b><u>Page</u></b>
2.1	Molecular weights, boiling points, and melting points of compounds investigated	7
2.2	Analyte percent recovery for the 50/50 (w/w) octadecylsilica/glass beads trap (1.0 g)	12
2.3	Analyte percent recovery for the 100% glass beads trap (1.5 g)	14
2.4	Analyte percent recovery for the 50/50 (w/w) Porapak Q®/glass beads trap (0.8 g)	15
3.1	Extraction and Trapping Conditions Used for Phases A-D	22
3.2	In-house Prepared Tablet Powder Mixture Uniformity With Liquid Extraction Method	26
3.3	SFE of MEVACOR® Tablets as a Function of Methanol-Modifier Percentage and Addition of Isopropylamine	38
3.4	SFE Reproducibility for Lovastatin (10 mg) from MEVACOR® Tablets with 15 % (1.0% (v/v) IPA) Methanol-Modified CO <sub>2</sub>	40
3.5	Traditional Liquid Extraction and SFE Solvent Use, Steps, and Time Comparisons	41
4.1	High/Low Extraction Parameters Investigated	49
4.2	Parameters Used for Fractional-Factorial Experiments	50
4.3	Least Square Coefficients, Reduced Fit, Response Percent Recovery	57
5.1	Phase I - Conditions for Ion-Pair/Supercritical Fluid Extraction Screening Study With Pure CO <sub>2</sub>	77
5.2	Extraction Conditions Used in Phases II-III	78

<b><u>Table</u></b>	<b><u>Description</u></b>	<b><u>Page</u></b>
5.3	Extraction Conditions and CO <sub>2</sub> Profile Used in Phase IV	80
6.1	Various Chromatographic Peak Parameters	133
6.2	Reproducibility Comparisons of Two ELSDs	135