

TABLE OF CONTENTS

ACKNOWLEDGEMENTS	v
LIST OF TABLES	viii
LIST OF FIGURES	x
CHAPTER ONE	
INTRODUCTION	1
CHAPTER TWO	
REVIEW OF LITERATURE	
Introduction	4
Development of Bud Necrosis	5
Vegetative Vigor	6
Shade	7
Fertilizer and Irrigation	9
Carbohydrate Status of Necrotic Buds	9
Gibberellin-Like Activity	10
Conclusion	11
Literature Cited	13
CHAPTER THREE	
ANATOMICAL DEVELOPMENTS IN THE ONSET OF GRAPEVINE BUD NECROSIS	
Abstract	15
Introduction	16
Materials and Methods	16
Time Frame of BN Development in ‘Riesling’ (1994)	16
Comparison of ‘Riesling’ and ‘Chardonnay’ (1995)	18
Results	18
Time Frame of BN Development in ‘Riesling’ (1994)	18
Comparison of ‘Riesling’ and ‘Chardonnay’ (1995)	29
Discussion	29
Conclusion	31
Literature Cited	32

TABLE OF CONTENTS (contd.)

CHAPTER FOUR

LOCALIZED CARBOHYDRATE AND ESSENTIAL NUTRIENT DEPRIVATION AS CAUSAL FACTORS IN GRAPEVINE BUD NECROSIS

Abstract	33
Introduction	34
Materials and Methods	36
Mineral Analysis	36
Carbohydrate Analysis	37
BN Assessment	38
Effect of Shade on BN in 'Riesling' Grapevines, 1994	38
Effect of Shade on BN in 'Riesling' Grapevines, 1996	40
Effect of Shoot Tipping on 'Riesling' BN, 1996	41
Mineral Nutrient and Carbohydrate Survey, 1995	41
Starch Rating of Grape Cultivars, 1996	42
Results	43
Shade Experiment, 1994	43
Shade Experiment, 1996	45
Shoot Tipping Experiment, 1996	55
Mineral Nutrient and Carbohydrate Survey, 1995	60
Starch Rating of Grape Cultivars, 1996	66
Discussion	66
Conclusion	75
Literature Cited	76

CHAPTER FIVE

CONCLUSIONS	79
-------------------	----

VITA	81
------------	----

LIST OF TABLES

Table 3.1.	Comparison of BN in Virginia ‘Riesling’ with Virginia ‘Chardonnay’ and New York ‘Riesling’	19
Table 4.1.	Key for Iodine-potassium iodide starch staining rating	44
Table 4.2.	Effect of different timings of artificial shade on bud necrosis incidence in ‘Riesling’ at Somerset and Willowcroft Vineyards, 1994	46
Table 4.3.	Effect of node position on bud necrosis incidence in ‘Riesling’ grapevines at Somerset and Willowcroft Vineyards, 1994	47
Table 4.4.	Total nonstructural carbohydrates level in ‘Riesling’ grapevines at Somerset Vineyard, 1994	48
Table 4.5.	Total nonstructural carbohydrates level in ‘Riesling’ grapevines at Willowcroft Vineyard, 1994	49
Table 4.6.	Effect of shade at different timings on BN in ‘Riesling’ during 1996	51
Table 4.7.	Carbohydrate analysis of shaded and unshaded ‘Riesling’ vines at Willowcroft Vineyard, Virginia during 1996	53
Table 4.8.	BN and fruit set measurements of shoot-tipped and non-shoot-tipped ‘Riesling’ vines during 1996	56
Table 4.9.	Carbohydrate analysis of shoot-tipped and non-shoot-tipped ‘Riesling’ vines at AHS AREC, Winchester, VA, 1996	57
Table 4.10.	Concentrations of mineral elements in ‘Riesling’ and ‘Chardonnay’ cultivars sampled from Prince Michel Vineyard during 1995	61
Table 4.11.	Carbohydrate level in bud tissues of ‘Riesling’ and ‘Chardonnay’ of Prince Michel and AHS AREC Vineyards during 1995	63
Table 4.12.	Carbohydrate level in leaf tissues of ‘Riesling’ and ‘Chardonnay’ of Prince Michel and AHS AREC Vineyards during 1995	64

LIST OF TABLES (contd.)

Table 4.13.	Carbohydrate level in stem tissues of 'Riesling' and 'Chardonnay' of Prince Michel and AHS AREC Vineyards during 1995.....	65
Table 4.14.	Effect of node position on starch deposits and BN in bud tissues of grape cultivars 'Riesling', 'Viognier', 'Syrah', and 'Chardonnay' at AHS AREC during 1996...	69

List of Figures

Fig. 3.1.	Healthy ‘Riesling’ buds without any cell abnormalities at 46 and 56 days after budbreak.....	21
Fig. 3.2.	Appearance of distorted and compressed zone of cells at 66 days after budbreak in ‘Riesling’ buds from Somerset Vineyard, Virginia	22
Fig. 3.3.	Zone of compression is visible at the base of the primary axis at the junction of the nodal cushion where the cells have undergone secondary thickening and the younger cells at the base of the primary axis	23
Fig. 3.4.	Three to four distorted cell layers were observed at 76 days after budbreak in ‘Riesling’ grapevines of Somerset Vineyard, Virginia.....	24
Fig. 3.5.	Seven to eight distorted layers of cells are visible at 86 and 96 days after budbreak in ‘Riesling’ vines	25
Fig. 3.6.	Localized groups of distorted cells in Willowcroft ‘Riesling’ buds at 66 days after budbreak.....	26
Fig. 3.7.	SEM micrograph showing a zone of compression at the base of the primary axis and extending to the base of the secondary buds and showing compressed layers of cells at the base of the primary axis observed at 76 days after budbreak.....	27
Fig. 3.8.	SEM micrograph of a bud where the primary axis and the secondary axis have separated from the nodal cushion.....	28
Fig. 4.1.	Relationship between shoot diameter (measured at the widest point of the second basal internode) and bud necrosis incidence in ‘Riesling’ at Somerset Vineyard	50
Fig. 4.2.	Effect of node position in shaded and non-shaded ‘Riesling’ vines at Willowcroft Vineyard during 1996	52
Fig. 4.3.	Effect of node position on BN in shoot-tipped and non-shoot-tipped ‘Riesling’ vines during 1996.....	59
Fig. 4.4.	Comparison of starch levels from 50 to 80 days after budbreak within each grape cultivar at AHS AREC during 1996	67

List of Figures (contd.)

Fig. 4.5.	Comparison of starch levels among four grape cultivars within each sampling date at AHS AREC during 1996.....	68
Fig. 4.6.	Correlation of BN with starch deposits in bud tissues of 'Riesling', 'Viognier', 'Syrah', and 'Chardonnay' at AHS AREC.....	70